



**Figure 1**




**Figure 1 (con't)**



**C**



UNIVERSITY OF CALIFORNIA  
SAN DIEGO




**Figure 2**






**Figure 3**





**Figure 4**



**GFP**

 **mt**

**mt = KISVYLTAAVVG****FVAYGILKWYRGT**

+

+

+

RECEIVED  
JAN 10 1992

10000447.070902



**Figure 5**


				domain is a mitochondrial targeting signal	domain is required for function
M11L	K	ISVYLTAADVGFVAYGIL	K WYRGT	Y	Y
Bcl-2	K	TLLSLALVGACITLGAYLS	K K	Y	Y/N
Bcl-X <sub>L</sub>	K	WFLTGMTVAGVLLGSLFS	K K	Y	Y/N
Boo/Diva	K	LLIQAFLSGFFATAIFFIW	K RL	?	?
CED-9	K	WSMIGAGVTAGAIGIVGVVCG	K MMFSLK	?	?
BHRF-1	K	FSWTLFLAGLTLSLLVICSYLFIS	K GRH	Y	Y
KSbcl-2	K	MTALLGSIALLATILAAVAMS	K R	?	?
Nip3	K	VFLPSLLLSHLLAIGLGIYIG	K RLTTSTSTF	Y	Y
Nix	K	VFIPSLFLSHVLALGLGIYIG	K RLSTPSA	Y	Y
	positive charge	18-24 aa putative membrane-spanning domain	positive charge    positive tail		

RECEIVED  
JAN 10 1991



**Figure 6**

**A**




000000000000  
100 000 000000

Figure 6 (con't)

**B**


Average percent difference in TMRE positive cells following apoptosis induction



GFP construct expressed by cells



Figure 7



RECEIVED  
JAN 11 1999  
FBI - NEW YORK

## Process to Identify M11L-interacting Proteins




Figure 8

10090440-070902



UNIVERSITY OF CALIFORNIA  
LIBRARY







Figure 9



016 JUN 25 1966 FBI LABORATORY

10099140 170000



Bak, a Bcl2 family member, was identified  
by Mass Spec

MASGQGGPPRQECGEPALPSASEEQVAQD

TEEVFRSYVFYRHHQQEQEAEGVAAPADPEM

VTLPLQPSSTMGQVGR**QLAIGDDINR**RYD

SEFQTMLQLQPTAENAYEYFTK**IATSLFE**

**SGINWGRVVALLGFGYR**LALHVVYQH

GLTGFLGQVTRFVDFMLHHCIARWIAQRGGWVA

ALNLGNPILNVLVLGWVLLGQFVVRFFKS

Figure 11



10040440.070902